



09/937839

1/29




2/29




09/937839

3/29



09/937839


4/29



09/937839

5/29

FIG. 5




09/937839

6/29

**FIG. 6**

Preparation of HVJ envelope vector




09/937839

7/29

FIG. 7

Protein profiles of HVJ, UV-inactivated HVJ,  
and HVJ envelope vector



09/937839

**FIG. 8**

8/29

Electron micrograph of an HVJ envelope vector

(1) Untreated HVJ



100 nm

(2) HVJ containing no DNA, which was subjected to an octylglucoside treatment



(3) HVJ containing DNA, which was subjected to an octylglucoside treatment



201220-688660





09/937839

**FIG. 9A**

9/29

Effects of octylglucoside on gene transfer  
by HVJ envelope vector


**FIG. 9B**

09/937839

10/29

FIG. 9C

Effects of octylglucoside on gene transfer  
by HVJ envelope vector




09/937839

FIG. 10A

11/29

Effects of protamine sulfate on gene  
transfer by HVJ envelope vector




00/937839

12/29

**FIG. 10B**

Effects of protamine sulfate on gene transfer by HVJ envelope vector





09/937839

13/29

**FIG. 11A**


Effects of DNA amounts on gene expression using frozen HVJ envelope which has been treated with octylglucoside



14/29

**FIG. 11B**

Effects of DNA amounts on gene expression  
by HVJ envelope vector




09/937839

15/29

FIG.12

Effects of HVJ titer on gene expression





09/937839

16/29

**FIG. 13A**



Study of irradiation amount in UV inactivation

**FIG. 13B**Study of treatment concentration  
in SPL inactivation



09/937839

17/29


**FIG. 14****SAS: Effects of infection time and protamine sulfate concentration****FIG. 15****HAEC: Effects of infection time and protamine sulfate concentration**

09/937839

18/29

**FIG. 16A**

Luciferase activity by HVJ envelope —  
vector in mouse liver




09/937839

19/29

**FIG. 16B**

Luciferase activity by HVJ envelope vector  
in mouse uterus



09/937839

20/29

201220-6E84E66D

**FIG. 16C**

**LacZ expression by HVJ envelope vector in mouse uterus**



21/29

**FIG. 16D**

Gene transfer into rat brain  
 using new HVJ

**#1 HVJ-GFP**

HVJ-GFP of 10,000 HAU was administered to SD rats (male, body weight: 300 to 400 g) via the cisterna magna or via the carotid artery. Samples were taken three to four days later.

Live sections were prepared, which were subjected to observation under fluorescence microscopy.

**(administration via the cisterna magna) ①**



Incorporation into the brain surface was confirmed.

No incorporation into deep portions of the brain was confirmed.

No incorporation into the choroid plexus was confirmed, either. → administration via the cisterna magna is considered to result in permeation through the intrathecal space, so that expression is usually observed in the choroid plexus

**(administration via the carotid artery) ②, ③**


Significant expression was confirmed on the administered side (left side). Expression was confirmed not only in the brain surface portions but also in the basal ganglia portion. Expression was also confirmed in the brain surface of the other brain, which was considered to have resulted from a flow to the other side through a collateral flow.

**② Administration via the carotid artery****③ Administration via the carotid artery****① Administration via the cisterna magna**

09/937839

22/29

**FIG. 16E** Inhibition of VEGF-induced angiogenesis  
by gene transfer using HVJ



09/937839

23/29



FIG. 16F


201220" 6E82E660/6

Introduction of FITC-ODN into BHK-21 cells by HVJ envelope vector

FIG. 17A

60 min

Fluorescence image



Phase-contrast image





FIG. 17B


10 min





09/937839

25/29


**FIG. 18A**Gene transfer into NALM-6 by  
centrifugation with HVJ envelope

Protamine sulfate concentration and centrifugation

201220" 6E8/E660

09/937839

26/29


**FIG. 18B**Gene transfer into CCRF-CEM by:  
centrifugation with HVJ envelope

09/937839

27/29

**FIG. 18C**Gene transfer into K-562 by centrifugation  
with HVJ envelope

(15 K rpm, 10 min, 20°C)




09/937839

28/29

**FIG. 19**

Gene transfer into mouse melanoma (B16-F1)  
mass using HVJ envelope



09/937839

29/29

FIG. 20




FIG. 21

